

Localization on short-range potentials in dissipative quantum mechanics

Akakii Melikidze

Physics Department, Princeton University, Princeton, NJ 08544

(February 1, 2008)

In this article the problem of the existence of a state localized on a weak short-range attractive potential in the presence of dissipation is considered. It is shown that, contrary to the pure quantum case, a localized state is produced in any number of dimensions, while in low dimensions dissipation leads to much stronger localization. The results have physical implications for the dissipative dynamics of objects such as heavy particles in Fermi liquids and for superconductivity in high- T_c materials.

Research in the theory of strongly interacting fermionic systems was rekindled soon after the discovery of high temperature cuprate superconductors when it was realized that Landau Fermi liquid theory can not describe the normal state of the underdoped materials. Since then, many ideas have been proposed to exhibit possible routes to non-Fermi liquid behavior. [1–6] While these theories differ in details, they all are intimately related to a peculiar property of fermionic liquids known as Anderson’s orthogonality catastrophe. [7] This property implies that the wave function of the liquid before and after an adiabatic insertion of an impurity has a vanishing overlap:

$$\langle \psi | \psi' \rangle \sim N^{-A}, \quad (1)$$

where N is the number of fermions in the liquid and A is the orthogonality exponent which can be expressed in terms of impurity scattering phase shifts. This result shows that interactions between fermions have singular effect and the fact that the non-interacting quasiparticles in Landau theory can often be defined is rather an exception than a principle.

Kondo [8] introduced a closely related overlap of the wave function of the liquid with an impurity at 0 and that with the impurity moved slowly to $\mathbf{x}$:

$$\langle \psi(0) | \psi(\mathbf{x}) \rangle \sim (\Lambda t)^{-K(\mathbf{x})}, \quad (2)$$

where t is a characteristic time of the impurity motion and Λ is a high energy cut-off. The exponent $K(\mathbf{x})$ is quite difficult to evaluate in general; however, it was shown by Schönhammer [9] that for small $\mathbf{x}$ ($\hbar = 1$ throughout the letter):

$$K(\mathbf{x}) \approx \frac{\eta \mathbf{x}^2}{2\pi}. \quad (3)$$

Here η is the friction coefficient that determines the viscous force $\mathbf{F} = -\eta \dot{\mathbf{x}}$ exerted on the moving impurity by the fermions in the liquid. That there is such a force can be easily seen by making a Galilean transformation from the reference frame of the liquid to that of the impurity. There, the fermionic liquid flows past the impurity and

the net force produced by scattering on the impurity has the same origin as the well-known residual resistance of metals. Moreover, this transformation allows a simple calculation of η which can be expressed in terms of the impurity’s transport scattering cross-section. [10]

The significance of Eq. (3) is that it relates an adiabatic effect — orthogonality catastrophe — with an essentially non-adiabatic effect — energy dissipation caused by the viscous force. In problems where the particle of interest can be singled out (like, for example, diffusion of heavy particles in Fermi liquids, the dynamics of topological defects, solitons, collective coordinates etc.) this connection provides a powerful tool that reduces the original strongly interacting problem to what is known as dissipative quantum mechanics of a single particle. [10] As for the Fermi liquid itself, one can not, in general, deduce the breakdown of Landau theory using these simple arguments. However, once the breakdown occurs, its direct cause — Anderson’s orthogonality catastrophe — can be linked to the dissipative dynamics of fermions. In the case of cuprate superconductors this idea is supported by experimental evidence of strongly damped quasiparticles. [11]

Motivated by these connections, I study the problem of localization of a particle on attractive short-range potentials in D dimensions in the presence of Ohmic dissipation at zero temperature. This problem is relevant to the localization of heavy particles on impurities in fermion liquids, pinning of vortices in superconductors [12] etc. It is also relevant to the problem of interaction between dissipative particles and to superconductivity. The question that I address is whether a short range attractive potential leads to localization of the particle and, if so, what are the properties of the localized state. I find that localization occurs for arbitrarily weak attractive potentials in any number of dimensions and for any strength of dissipation. This should be compared to the corresponding problem in the non-dissipative case [13] where it is known that weak potentials localize the particle in $D = 1$, lead to exponentially weak localization in $D = 2$ and do not lead to localization in $D \geq 3$. I suggest that

the much stronger localization in the dissipative case may be relevant to the physics of Anderson localization and the problem of high critical temperature in cuprate superconductors.

Consider a particle of mass m that moves in D dimensions in the presence of an attractive short-range potential $V(\mathbf{x}) = -\alpha\delta(\mathbf{x})$. The particle also experiences the action of a friction force that is linear in the particle's velocity: $\mathbf{F} = -\eta\dot{\mathbf{x}}$ (Ohmic case). The well-accepted microscopic approach to this problem is to think of the friction force as originating from coupling to an effective environment. The environmental degrees of freedom are eventually traced over to produce an effective description of the particle's dynamics. I use the imaginary (single) time path integral representation of the partition function. [14] This way, tracing over the environmental degrees of freedom reduces to path-integration over environmental variables, yielding the following effective action description: [15]

$$Z = \int \mathcal{D}\mathbf{x} e^{-S}, \quad (4)$$

$$S = \frac{T}{2} \sum_{n=-\infty}^{+\infty} (m\nu_n^2 + \eta|\nu_n|) |\mathbf{x}_n|^2 - \alpha \int_0^{1/T} d\tau \delta(\mathbf{x}(\tau)). \quad (5)$$

Here $\nu_n = 2\pi Tn$ are Matsubara frequencies: $\mathbf{x}(\tau) = \sum e^{i\nu_n\tau} \mathbf{x}_n$.

The $\eta|\nu_n|$ term in the effective action S is responsible for dissipation. Note that it has a non-analytic $|\nu_n|$ frequency dependence. This fact strongly complicates the problem because it leads to a non-local effective action when written in time representation. However, there are two limits in which the problem is somewhat simplified: the pure quantum limit and the dissipative limit. Indeed, the interplay between different terms in S produces a characteristic energy scale ω_0 . Then, in the pure quantum limit $\omega_0 \gg \eta/m$ the dissipative term in S is much smaller than the kinetic energy term and can, therefore, be neglected. The resulting pure quantum problem can be solved exactly. [13] In the opposite dissipative limit $\omega_0 \ll \eta/m$ the kinetic energy term can be neglected; its only purpose then is to set a high-energy cut-off.

To analyze the problem I employ the variational method. [14] This method exploits the fact that for an arbitrary trial action S_0 the following inequality puts an upper bound on the free energy:

$$F \leq F_0 + T\langle S - S_0 \rangle_{S_0}. \quad (6)$$

Here $F = T \ln Z = T \ln \int \mathcal{D}\mathbf{x} e^{-S}$, $F_0 = T \ln Z_0 = T \ln \int \mathcal{D}\mathbf{x} e^{-S_0}$ and $\langle S - S_0 \rangle_{S_0} = Z_0^{-1} \int \mathcal{D}\mathbf{x} (S - S_0) e^{-S_0}$. We are interested in the ground state energy: $E = \lim_{T \rightarrow 0} F$. Taking the low temperature limit in Eq. (6) we get:

$$E \leq E_0 + \lim_{T \rightarrow 0} T\langle S - S_0 \rangle_{S_0}. \quad (7)$$

Here $E_0 = \lim_{T \rightarrow 0} F_0$. I take for S_0 the action of an Ohmic-damped harmonic oscillator:

$$S_0 = \frac{T}{2} \sum_{n=-\infty}^{+\infty} (m\nu_n^2 + \eta|\nu_n| + m\omega_0^2) |\mathbf{x}_n|^2. \quad (8)$$

Here, the ω_0 of the oscillator will serve as a variational parameter. As we shall see below, all quantities that are necessary for the evaluation of Eq. (7) can be expressed through the expectation values of the operators that are diagonal in the x -representation. This means that we only need to know the diagonal elements of the equilibrium density matrix ρ_0 that corresponds to S_0 . However, the fact S_0 is quadratic implies that ρ_0 is Gaussian. Thus, in the $T \rightarrow 0$ limit we can write:

$$\rho_0(\mathbf{x}, \mathbf{x}) = \frac{1}{(2\pi\langle x^2 \rangle_0)^{\frac{D}{2}}} \exp \left\{ -\frac{\mathbf{x}^2}{2\langle x^2 \rangle_0} \right\}. \quad (9)$$

Here, $\langle x^2 \rangle_0 = \frac{1}{D} \langle \mathbf{x}^2 \rangle_0$ is the ground state expectation value of x^2 for the damped harmonic oscillator that corresponds to S_0 . This quantity can be easily evaluated:

$$\langle x^2 \rangle_0 = \lim_{T \rightarrow 0} T \sum_{n=-\infty}^{+\infty} \frac{1}{m\nu_n^2 + \eta|\nu_n| + m\omega_0^2}. \quad (10)$$

In the $T \rightarrow 0$ limit the sum can be substituted by an integral. Defining $r = 2m\omega_0/\eta$, we can write the resulting expression as:

$$\langle x^2 \rangle_0 = \frac{1}{\pi\eta} \times \begin{cases} \frac{1}{\sqrt{1-r^2}} \ln \frac{1+\sqrt{1-r^2}}{1-\sqrt{1-r^2}}, & r < 1; \\ \frac{2}{\sqrt{r^2-1}} \arctan \sqrt{r^2-1}, & r > 1. \end{cases} \quad (11)$$

In particular, in the dissipative regime $r \ll 1$ we get: $\langle x^2 \rangle_0 = (2/\pi\eta) \ln(1/r)$, while in the pure quantum regime $r \gg 1$ we obtain the correct quantum expression: $\langle x^2 \rangle_0 = 1/2m\omega_0$; at $r = 1$ $\langle x^2 \rangle_0$ is analytic and equals $2/\pi\eta = 1/\pi m\omega_0$. Note that $\langle x^2 \rangle_0$ for $r > 1$ can be obtained from the expression for $r < 1$ by an analytic continuation.

Next, using Eq. (9) we can evaluate:

$$\begin{aligned} \lim_{T \rightarrow 0} T\langle S - S_0 \rangle_0 &= \langle -\alpha\delta(\mathbf{x}) - \frac{m\omega_0^2}{2}\mathbf{x}^2 \rangle_0 \\ &= -\frac{\alpha}{(2\pi\langle x^2 \rangle_0)^{\frac{D}{2}}} - \frac{Dm\omega_0^2}{2} \langle x^2 \rangle_0. \end{aligned} \quad (12)$$

To evaluate E_0 we apply Hellmann-Feynman theorem: $\partial E_0 / \partial \omega_0 = m\omega_0 \langle x^2 \rangle_0$; this leads to:

$$E_0 = Dm \int_0^{\omega_0} d\omega'_0 \omega'_0 \langle x^2 \rangle_0. \quad (13)$$

Substituting Eqs. (12,13) into Eq. (7) and integrating by parts, we get a variational bound on the ground state energy:

$$E \leq -\frac{Dm}{2} \int_0^{\omega_0} d\omega'_0 \omega_0'^2 \frac{\partial}{\partial \omega'_0} \langle x^2 \rangle_0 - \frac{\alpha}{(2\pi \langle x^2 \rangle_0)^{\frac{D}{2}}}. \quad (14)$$

An immediate consequence of this result is the following statement: For any combination of non-zero values of dimensionality D , mass of the particle m and friction constant η the ground state energy $E < 0$. And since for any delocalized state $E \geq 0$ this implies that the particle is always localized! To see this, notice that for small enough values of ω_0 it follows from Eq. (11) that the first term in Eq. (14) is positive and $\propto \omega_0^2$ while the second term is negative and its absolute value is $\propto (\ln \frac{1}{\omega_0})^{-D/2}$. Therefore for sufficiently small ω_0 the second term dominates and the energy of the ground state is bounded from above by a negative value.

Minimizing Eq. (14) with respect to ω_0 we obtain an equation on ω_0 :

$$\frac{\alpha}{(2\pi)^{\frac{D}{2}}} = m\omega_0^2 \langle x^2 \rangle_0^{\frac{D}{2}+1}. \quad (15)$$

The upper bound on the ground state energy can be found numerically by solving Eqs. (14,15); this solution is presented below. However, in two limiting cases, $\omega_0 \ll \eta/m$ and $\omega_0 \gg \eta/m$, approximate analytical expressions can be obtained. These two limits are realized for weak and strong pinning potentials respectively. To be more precise, let us define α_c as the strength of the potential at which the minimizing value ω_0 satisfies $r = 2m\omega_0/\eta = 1$. Substituting this in Eqs. (11,15) we obtain:

$$\alpha_c = \frac{2^{D-1}}{\pi m} \eta^{1-\frac{D}{2}}. \quad (16)$$

For weak attractive potentials, such that $\alpha \ll \alpha_c$, we have the dissipative regime: $\omega_0 \ll \eta/m$. In this case, using the limiting form of Eq. (11) for $r \ll 1$ and solving Eqs. (14,15) we obtain:

$$E_{\text{diss}} \leq -\frac{\eta}{2\pi m} \left(\frac{\alpha}{\alpha_c} \right) \left(\ln \frac{\alpha_c}{\alpha} \right)^{-\frac{D}{2}}. \quad (17)$$

This result shows that the localization on sufficiently weak attractive potentials is dominated by dissipative dynamics and is much stronger compared to the non-dissipative case. Indeed, note the linear dependence of E_{diss} on α in Eq. (17) vs. exact results in the pure quantum case: [13] $E_{\text{quant}} = -m\alpha^2/2$ in $D = 1$; $E_{\text{quant}} = -(m\lambda^2)^{-1} \exp(-2\pi/m\alpha)$ in $D = 2$ (λ is the short-distance cut-off); no localization on weak potentials in $D \geq 3$.

In the opposite limit of sufficiently strong potentials, such that $\alpha \gg \alpha_c$, we have the pure quantum regime: $\omega_0 \gg \eta/m$. Here the situation depends on the dimensionality. In $D = 1$, using the limiting form of Eq. (11) for $r \gg 1$ and solving Eqs. (14,15), we obtain:




FIG. 1. Numerical minimization of Eq. (14) (diamonds) shows the dependence of the upper bound on the ground state energy E (in units of η/m) on the strength of the potential α/α_c for $D = 1$. Also shown are E_{diss} , Eq. (17) (dotted line) and E_{quant} , Eq. (18) (dashed line).

$$E_{\text{quant}}^{(D=1)} \leq -\frac{m\alpha^2}{\pi}. \quad (18)$$

This bound is reasonably close to the exact quantum-mechanical result $E = -m\alpha^2/2$. In dimensions $D \geq 2$ the pure quantum problem of a strong ($\alpha \gg \alpha_c$) attractive potential acquires a short distance cut-off (bandwidth) dependence; this is a direct consequence of the fact that δ -function is poorly defined in $D > 1$. However, it should be emphasized that the particle is always localized in any D (see the arguments above)! While the question about the ground state energy in the strong potential case lies outside the main line of this letter, it could be mentioned that, in the first approximation, $E(\alpha \gg \alpha_c)$ is bounded from above by $E_{\text{diss}}(\alpha \sim \alpha_c) \sim -\eta/m$ for any D .

Finally, let us discuss the numerical minimization of Eq. (14). The results of the solution for $D = 1$ are presented in Fig. 1. A crossover between the $\propto \alpha^2$ behavior in the pure quantum regime (large α) and the $\propto \alpha$ behavior in the dissipative regime (small α) occurs at $\alpha \sim \alpha_c$. Note the much stronger localization that occurs for weak potentials compared to the pure quantum case. One can see that the two limiting expressions, Eqs. (17,18) describe the numerical results well in their respective limits.

To understand the qualitatively different localization in the dissipative case let us first recall how localization occurs in the non-dissipative case. There, it is determined by the competition of two tendencies of the particle: to localize and thus lower its potential energy and to diffuse away thus lowering its kinetic energy. In the dissipative case, however, the diffusion is strongly suppressed by the Anderson's orthogonality catastrophe (see Eq. (2)). Indeed, as was shown by Hakim and Ambegaokar [16], the mean-squared displacement of a free Ohmic-damped particle at $T = 0$ (this is known as quan-

tum Brownian motion problem) grows very slowly with time:

$$\langle x^2 \rangle = \frac{2}{\pi\eta} \ln \frac{\eta t}{m}. \quad (19)$$

(This can also be shown [10] to be a direct consequence of the fluctuation-dissipation theorem of Callen and Welton [17]). Thus, strictly speaking, the diffusion coefficient of the Ohmic-damped particle is zero. Therefore, the tendency to lower potential energy by localization dominates over diffusion and leads to localization for any D , in contrast to the pure quantum case. Meanwhile, in one dimension dissipation leads to much stronger localization on weak potentials compared to the non-dissipative case.

Physical implications of these results are quite interesting. For example, if one considers the problem of diffusion of a heavy particle in fermionic liquids with impurities one arrives at the conclusion that the theory of Anderson localization breaks down due to the presence of dissipation (orthogonality catastrophe). In particular, no metal-insulator transition is possible for heavy particles in $D = 3$ since in any dimensionality all states are localized.

Perhaps, a more important implication is for the interaction between damped particles. The well-known effect of attractive short-range interactions is superconductivity. In conventional superconductors superconductivity is known to arise from Cooper pair formation by electrons near the Fermi surface. For such electrons the density of states can be taken constant. Therefore, the problem of Cooper pairing becomes analogous (after a transformation to the center-of-mass and relative coordinates) to single-particle localization on attractive short-range potentials in $D = 2$ where a single particle has constant density of states. In the latter problem, the binding energy is exponentially small: [13] $E \sim W \exp(-1/\alpha\nu)$, where W is the band-width, α is the strength of the potential and $\nu = m/2\pi$ is the single-particle density of states in $D = 2$. This explains why the critical temperature T_c is so low in conventional superconductors. While no single accepted theory of superconductivity in the high- T_c cuprate superconductors exists yet, most attempts to explain high T_c begin by considering strong interactions: $\alpha\nu \sim 1$, while still following the BCS ideas (extended to the strong-coupling case by Eliashberg). Here, I suggest that in systems where the Landau Fermi liquid theory breaks down and the dynamics of fermions is dissipative (governed by the orthogonality catastrophe) the nature of Cooper pairing changes *qualitatively*. Most importantly, the binding energy becomes much bigger and is essentially linear in α for weak attractive interactions (see Eq. (17)). This means that any weak second-order (fluctuation-induced) process is, in principle, capable of producing sufficiently high T_c .

It should be emphasized, however, that this model of superconductivity is a phenomenological one since the

arguments given in the beginning of the letter can not be used to deduce the breakdown of the Landau theory. However, if the breakdown does occur, this model is believed to capture the qualitative features of the physical picture.

I would like to thank F. D. M. Haldane for sharing his insight into the problem. I have also benefited from discussions with L. P. Gor'kov and O. Motrunich. This work was supported by NSF MRSEC DMR-9809483.

-
- [1] P. W. Anderson, *The theory of superconductivity in the high- T_c cuprates* (Princeton University Press, Princeton, 1997).
 - [2] C. M. Varma, Phys. Rev. Lett. **63**, 1996 (1989); C. M. Varma, Z. Nussinov, W. van Saarloos, cond-mat/0103393, submitted to Physics Reports.
 - [3] P. A. Lee and N. Nagaosa, Phys. Rev. B **46**, 5621 (1992); P. A. Lee, Physica C **317**, 194 (1999).
 - [4] D. J. Scalapino, Phys. Rep. **250**, 329 (1995); D. Pines, Z. Phys. B **103**, 129 (1997); T. Moriya and K. Ueda, Adv. Phys. **49**, 555 (2000). Ar. Abanov, A. V. Chubukov and J. Schmalian, cond-mat/0010403 (unpublished).
 - [5] S. Sachdev, Science **288**, 475 (2000).
 - [6] A. J. Schofield, Contemp. Phys. **40**, 95 (1999).
 - [7] P. W. Anderson, Phys. Rev. Lett. **18**, 1049 (1967).
 - [8] J. Kondo, in *Fermi surface effects*, Springer Series in Solid State Physics, vol. 77, eds. J. Kondo and A. Yoshimori (Springer, Berlin, 1988).
 - [9] K. Schönhammer, Phys. Rev. B **43**, 11323 (1991).
 - [10] U. Weiss, *Quantum dissipative systems* (World Scientific, Singapore, 1993).
 - [11] J. Orenstein and A. J. Millis, Science **288**, 468 (2000).
 - [12] A. Melikidze, to appear in Phys. Rev. B.
 - [13] L. D. Landau and E. M. Lifshits, *Quantum mechanics: non-relativistic theory* (Pergamon, London, 1958).
 - [14] R. P. Feynman, *Statistical Mechanics* (Benjamin, New York, 1972).
 - [15] A. O. Caldeira and A. J. Leggett, Phys. Rev. Lett. **46**, 211 (1981); Ann. Phys. **149**, 374 (1983); Physica **121A**, 587 (1983).
 - [16] V. Hakim and V. Ambegaokar, Phys. Rev. A **32**, 423 (1985).
 - [17] H. B. Callen and T. A. Welton, Phys. Rev. **83**, 34 (1951).